

Refid **Bibliograph**

- 5 M. Guo, A. C. Baruch, E. G. Silva, Y. J. Jan, E. Lin, N. Sneige and M. T. Deavers. Efficacy of p16 and ProExC immunostaining in the detection of high-grade cervical intraepithelial neoplasia and cervical carcinoma. *Am J Clin Pathol.* 2011. 135:212-20
- 14 E. C. Pirog, K. D. Quint and R. K. Yantiss. P16/CDKN2A and Ki-67 enhance the detection of anal intraepithelial neoplasia and condyloma and correlate with human papillomavirus detection by polymerase chain reaction. *Am J Surg Pathol.* 2010. 34:1449-55
- 36 Y. S. Choi, W. D. Kang, S. M. Kim, Y. D. Choi, J. H. Nam, C. S. Park and H. S. Choi. Human papillomavirus L1 capsid protein and human papillomavirus type 16 as prognostic markers in cervical intraepithelial neoplasia 1. *Int J Gynecol Cancer.* 2010. 20:288-93
- 39 A. Kreuter, M. Jesse, A. Potthoff, N. H. Brockmeyer, T. Gambichler, M. Stucker, F. G. Bechara, H. Pfister and U. Wieland. Expression of proliferative biomarkers in anal intraepithelial neoplasia of HIV-positive men. *J Am Acad Dermatol.* 2010. 63:490-8
- 62 S. M. Bean, R. S. Meara, R. T. Vollmer, M. G. Conner, D. R. Crowe, L. Novak, I. A. Eltoum, S. J. Robboy and D. C. Chhieng. p16 Improves interobserver agreement in diagnosis of anal intraepithelial neoplasia. *J Low Genit Tract Dis.* 2009. 13:145-53
- 68 I. Tsoumpou, M. Arbyn, M. Kyrgiou, N. Wentzensen, G. Koliopoulos, P. Martin-Hirsch, V. Malamou-Mitsi and E. Paraskevaidis. p16(INK4a) immunostaining in cytological and histological specimens from the uterine cervix: a systematic review and meta-analysis. *Cancer Treat Rev.* 2009. 35:210-20

- 97 J. Monsonego, G. Pollini, M. J. Evrard, P. Sednaoui, L. Monfort, D. Quinzat, R. Dachez and K. Syrjanen. Linear array genotyping and hybrid capture II assay in detecting human papillomavirus genotypes in women referred for colposcopy due to abnormal Papanicolaou smear. *Int J STD AIDS*. 2008. 19:385-92
- 102 A. P. Pinto, N. F. Schlecht, T. Y. Woo, C. P. Crum and E. S. Cibas. Biomarker (ProEx C, p16(INK4A), and MiB-1) distinction of high-grade squamous intraepithelial lesion from its mimics. *Mod Pathol*. 2008. 21:1067-74
- 112 J. E. Bernard, M. O. Butler, L. Sandweiss and N. Weidner. Anal intraepithelial neoplasia: correlation of grade with p16INK4a immunohistochemistry and HPV in situ hybridization. *Appl Immunohistochem Mol Morphol*. 2008. 16:215-20
- 132 A. A. El-Ghobashy, A. M. Shaaban, J. Innes, W. Prime and C. S. Herrington. Differential expression of cyclin-dependent kinase inhibitors and apoptosis-related proteins in endocervical lesions. *Eur J Cancer*. 2007. 43:2011-8
- 148 G. R. Focchi, I. D. Silva, N. C. Nogueira-de-Souza, C. Dobo, C. T. Oshima and J. N. Stavale. Immunohistochemical expression of p16(INK4A) in normal uterine cervix, nonneoplastic epithelial lesions, and low-grade squamous intraepithelial lesions. *J Low Genit Tract Dis*. 2007. 11:98-104
- 149 S. M. Bean, I. Eltoun, D. K. Horton, L. Whitlow and D. C. Chhieng. Immunohistochemical expression of p16 and Ki-67 correlates with degree of anal intraepithelial neoplasia. *Am J Surg Pathol*. 2007. 31:555-61

- 153 M. Hampl, N. Wentzensen, S. Vinokurova, M. von Knebel-Doeberitz, C. Poremba, H. G. Bender and V. Kueppers. Comprehensive analysis of 130 multicentric intraepithelial female lower genital tract lesions by HPV typing and p16 expression profile. *J Cancer Res Clin Oncol.* 2007. 133:235-45
- 177 A. E. Walts, J. Lechago and S. Bose. P16 and Ki67 immunostaining is a useful adjunct in the assessment of biopsies for HPV-associated anal intraepithelial neoplasia. *Am J Surg Pathol.* 2006. 30:795-801
- 189 M. Benevolo, M. Mottolese, F. Marandino, G. Vocaturo, R. Sindico, G. Piperno, L. Mariani, I. Sperduti, P. Canalini, R. P. Donnorso and A. Vocaturo. Immunohistochemical expression of p16(INK4a) is predictive of HR-HPV infection in cervical low-grade lesions. *Mod Pathol.* 2006. 19:384-91
- 242 S. S. Wang, M. Trunk, M. Schiffman, R. Herrero, M. E. Sherman, R. D. Burk, A. Hildesheim, M. C. Bratti, T. Wright, A. C. Rodriguez, S. Chen, A. Reichert, C. von Knebel Doeberitz, R. Ridder and M. von Knebel Doeberitz. Validation of p16INK4a as a marker of oncogenic human papillomavirus infection in cervical biopsies from a population-based cohort in Costa Rica. *Cancer Epidemiol Biomarkers Prev.* 2004. 13:1355-60
- 589 M. Benevolo, I. Terrenato, M. Mottolese, F. Marandino, P. Muti, M. Carosi, F. Rollo, L. Ronchetti, L. Mariani, G. Vocaturo and A. Vocaturo. Comparative evaluation of nm23 and p16 expression as biomarkers of high-risk human papillomavirus infection and cervical intraepithelial neoplasia 2(+) lesions of the uterine cervix. *Histopathology.* 2010. 57:580-6
- 593 W. J. Zeng, Y. Li, H. L. Fei, X. D. Cheng, F. Ye, W. G. Lu and X. Xie. The value of p16ink4a expression by fluorescence in situ hybridization in triage for high risk HPV positive in cervical cancer screening. *Gynecol Oncol.* 2011. 120:84-8

- 594 M. G. Dijkstra, D. A. Heideman, S. C. de Roy, L. Rozendaal, J. Berkhof, K. van Krimpen, K. van Groningen, P. J. Snijders, C. J. Meijer and F. J. van Kemenade. p16(INK4a) immunostaining as an alternative to histology review for reliable grading of cervical intraepithelial lesions. *J Clin Pathol.* 2010. 63:972-7
- 622 J. Hanprasertpong, K. Tungsinmunkong, S. Chichareon, V. Wootipoom, A. Geater, R. Buhachat and S. Boonyapipat. Correlation of p53 and Ki-67 (MIB-1) expressions with clinicopathological features and prognosis of early stage cervical squamous cell carcinomas. *J Obstet Gynaecol Res.* 2010. 36:572-80
- 639 S. Hovland, S. Muller, H. Skomedal, M. Mints, J. Bergstrom, K. L. Wallin, F. Karlsen, B. Johansson and S. Andersson. E6/E7 mRNA expression analysis: a test for the objective assessment of cervical adenocarcinoma in clinical prognostic procedure. *Int J Oncol.* 2010. 36:1533-9
- 663 C. Bergeron, J. Ordi, D. Schmidt, M. J. Trunk, T. Keller and R. Ridder. Conjunctive p16INK4a testing significantly increases accuracy in diagnosing high-grade cervical intraepithelial neoplasia. *Am J Clin Pathol.* 2010. 133:395-406
- 666 M. Mimica, S. Tomic, G. Kardum, I. D. Hofman, V. Kaliterna and L. Pejkovic. Ki-67 quantitative evaluation as a marker of cervical intraepithelial neoplasia and human papillomavirus infection. *Int J Gynecol Cancer.* 2010. 20:116-9
- 669 G. C. Tan, S. Norlatiffah, N. A. Sharifah, G. Razmin, M. S. Shiran, A. Z. Hatta and H. O. Paul-Ng. Immunohistochemical study of p16 INK4A and survivin expressions in cervical squamous neoplasm. *Indian J Pathol Microbiol.* 2010. 53:1-6

- 683 W. Sui, M. Ou, Y. Dai, J. Chen, H. Lan, Q. Yan and H. Huang. Gain of the human telomerase RNA gene TERC at 3q26 is strongly associated with cervical intraepithelial neoplasia and carcinoma. *Int J Gynecol Cancer*. 2009. 19:1303-6
- 689 M. Z. Huang, H. B. Li, X. M. Nie, X. Y. Wu and X. M. Jiang. An analysis on the combination expression of HPV L1 capsid protein and p16INK4a in cervical lesions. *Diagn Cytopathol*. 2010. 38:573-8
- 691 I. Takac, D. Arko, T. Kodric, M. Poljak, A. Zagorac, A. Erjavec-Skerget and N. Kokalj-Vokac. Human telomerase gene amplification and high-risk human papillomavirus infection in women with cervical intra-epithelial neoplasia. *J Int Med Res*. 2009. 37:1588-95
- 721 P. Conesa-Zamora, A. Domenech-Peris, F. J. Orantes-Casado, S. Ortiz-Reina, L. Sahuquillo-Frias, J. Acosta-Ortega, J. Garcia-Solano and M. Perez-Guillermo. Effect of human papillomavirus on cell cycle-related proteins p16, Ki-67, Cyclin D1, p53, and ProEx C in precursor lesions of cervical carcinoma: a tissue microarray study. *Am J Clin Pathol*. 2009. 132:378-90
- 723 M. del Pino, S. Garcia, V. Fuste, I. Alonso, P. Fuste, A. Torne and J. Ordi. Value of p16(INK4a) as a marker of progression/regression in cervical intraepithelial neoplasia grade 1. *Am J Obstet Gynecol*. 2009. 201:488 e1-7
- 757 Z. Tu, A. Zhang, R. Wu, J. Jiang, Y. Li, N. Wulan, J. Li, Y. Zhang, Z. Chen and L. Wei. Genomic amplification of the human telomerase RNA gene for differential diagnosis of cervical disorders. *Cancer Genet Cytogenet*. 2009. 191:10-6

- 770 A. E. Walts and S. Bose. p16, Ki-67, and BD ProExC immunostaining: a practical approach for diagnosis of cervical intraepithelial neoplasia. *Hum Pathol.* 2009. 40:957-64
- 784 D. Aximu, A. Azad, R. Ni, T. Colgan and S. Nanji. A pilot evaluation of a novel immunohistochemical assay for topoisomerase II-alpha and minichromosome maintenance protein 2 expression (ProEx C) in cervical adenocarcinoma in situ, adenocarcinoma, and benign glandular mimics. *Int J Gynecol Pathol.* 2009. 28:114-9
- 798 J. Ordi, S. Garcia, M. del Pino, S. Landolfi, J. Alonso, L. Quinto and A. Torne. P16 INK4a immunostaining identifies occult CIN lesions in HPV-positive women. *Int J Gynecol Pathol.* 2009.
- 830 C. M. Gurrola-Diaz, A. E. Suarez-Rincon, G. Vazquez-Camacho, G. Buonocunto-Vazquez, S. Rosales-Quintana, N. Wentzensen and M. von Knebel Doeberitz. P16INK4a immunohistochemistry improves the reproducibility of the histological diagnosis of cervical intraepithelial neoplasia in cone biopsies. *Gynecol Oncol.* 2008. 111:120-4
- 865 R. E. Badr, A. E. Walts, F. Chung and S. Bose. BD ProEx C: a sensitive and specific marker of HPV-associated squamous lesions of the cervix. *Am J Surg Pathol.* 2008. 32:899-906
- 878 M. Branca, M. Ciotti, C. Giorgi, D. Santini, L. Di Bonito, S. Costa, A. Benedetto, D. Bonifacio, P. Di Bonito, P. Paba, L. Accardi, S. Syrjanen, C. Favalli and K. Syrjanen. Predicting high-risk human papillomavirus infection, progression of cervical intraepithelial neoplasia, and prognosis of cervical cancer with a panel of 13 biomarkers tested in multivariate modeling. *Int J Gynecol Pathol.* 2008. 27:265-73

- 890 L. C. Horn, A. Reichert, A. Oster, S. F. Arndal, M. J. Trunk, R. Ridder, O. F. Rassmussen, K. Bjelkenkrantz, P. Christiansen, M. Eck, T. Lorey, V. R. Skovlund, T. Ruediger, V. Schneider and D. Schmidt. Immunostaining for p16INK4a used as a conjunctive tool improves interobserver agreement of the histologic diagnosis of cervical intraepithelial neoplasia. *Am J Surg Pathol.* 2008. 32:502-12
- 893 S. Muller, C. Flores-Staino, B. Skyldberg, A. C. Hellstrom, B. Johansson, B. Hagmar, K. L. Wallin and S. Andersson. Expression of p16INK4a and MIB-1 in relation to histopathology and HPV types in cervical adenocarcinoma. *Int J Oncol.* 2008. 32:333-40
- 956 J. Hariri and A. Oster. The negative predictive value of p16INK4a to assess the outcome of cervical intraepithelial neoplasia 1 in the uterine cervix. *Int J Gynecol Pathol.* 2007. 26:223-8
- 959 A. Sharma, M. Rajappa, A. Saxena and M. Sharma. Telomerase activity as a tumor marker in Indian women with cervical intraepithelial neoplasia and cervical cancer. *Mol Diagn Ther.* 2007. 11:193-201
- 964 J. Shi, H. Liu, M. Wilkerson, Y. Huang, S. Meschter, W. Dupree, C. Schuerch and F. Lin. Evaluation of p16INK4a, minichromosome maintenance protein 2, DNA topoisomerase IIalpha, ProEX C, and p16INK4a/ProEX C in cervical squamous intraepithelial lesions. *Hum Pathol.* 2007. 38:1335-44
- 965 M. I. Rosa, L. R. Medeiros, M. C. Bozzetti, J. Fachel, E. Wendland, R. R. Zanini, A. B. Moraes and D. D. Rosa. Accuracy of telomerase in cervical lesions: a systematic review. *Int J Gynecol Cancer.* 2007. 17:1205-14

- 984 C. G. Hsu, P. H. Wang, J. L. Ko, G. D. Chen, H. Chang, S. F. Yang, S. C. Chen and L. Y. Lin. Concurrent high expression of human telomerase reverse transcriptase and human nonmetastatic clone 23 in high-grade squamous intraepithelial neoplasia and squamous cell carcinoma of uterine cervix. *Int J Gynecol Cancer*. 2007. 17:851-7
- 1041 M. Branca, C. Giorgi, M. Ciotti, D. Santini, L. Di Bonito, S. Costa, A. Benedetto, D. Bonifacio, P. Di Bonito, P. Paba, L. Accardi, L. Mariani, M. Ruutu, S. Syrjanen, C. Favalli and K. Syrjanen. Upregulation of telomerase (hTERT) is related to the grade of cervical intraepithelial neoplasia, but is not an independent predictor of high-risk human papillomavirus, virus persistence, or disease outcome in cervical cancer. *Diagn Cytopathol*. 2006. 34:739-48
- 1045 M. Branca, C. Giorgi, M. Ciotti, D. Santini, L. Di Bonito, S. Costa, A. Benedetto, D. Bonifacio, P. Di Bonito, P. Paba, L. Accardi, L. Mariani, M. Ruutu, S. M. Syrjanen, C. Favalli and K. Syrjanen. Over-expression of topoisomerase IIalpha is related to the grade of cervical intraepithelial neoplasia (CIN) and high-risk human papillomavirus (HPV), but does not predict prognosis in cervical cancer or HPV clearance after cone treatment. *Int J Gynecol Pathol*. 2006. 25:383-92
- 1060 K. Pavlakis, I. Messini, S. Athanassiadou, E. Kyrodinou, A. Pandazopoulou, T. Vrekoussis and E. N. Stathopoulos. Endocervical glandular lesions: a diagnostic approach combining a semi-quantitative scoring method to the expression of CEA, MIB-1 and p16. *Gynecol Oncol*. 2006. 103:971-6
- 1068 R. Santopietro, I. Shabalova, N. Petrovichev, V. Kozachenko, T. Zakharova, J. Pajanidi, J. Podistov, G. Chemeris, L. Sozaeva, E. Lipova, I. Tsidaeva, O. Ivanchenko, A. Pshepurko, S. Zakharenko, R. Nerovjna, L. Kljukina, O. Erokhina, M. Branovskaja, M. Nikitina, V. Grunberga, A. Grunberg, A. Juschenko, M. Cintonino, P. Tosi, K. Syrjanen and S. Syrjanen. Cell cycle regulators p105, p107, Rb2/p130, E2F4, p21CIP1/WAF1, cyclin A in predicting cervical intraepithelial neoplasia, high-risk human papillomavirus
- 1092 B. Samama, D. Lipsker and N. Boehm. p16 expression in relation to human papillomavirus in anogenital lesions. *Hum Pathol*. 2006. 37:513-9

- 1121 M. Ishikawa, T. Fujii, M. Saito, I. Nindl, A. Ono, K. Kubushiro, K. Tsukazaki, M. Mukai and S. Nozawa. Overexpression of p16 INK4a as an indicator for human papillomavirus oncogenic activity in cervical squamous neoplasia. *Int J Gynecol Cancer*. 2006. 16:347-53
- 1151 M. Lorenzato, S. Caudroy, C. Bronner, G. Evrard, M. Simon, A. Durlach, P. Birembaut and C. Clavel. Cell cycle and/or proliferation markers: what is the best method to discriminate cervical high-grade lesions?. *Hum Pathol*. 2005. 36:1101-7
- 1183 M. Branca, C. Giorgi, D. Santini, L. Di Bonito, M. Ciotti, S. Costa, A. Benedetto, E. A. Casolati, C. Favalli, P. Paba, P. Di Bonito, L. Mariani, S. Syrjanen, D. Bonifacio, L. Accardi, F. Zanconati and K. Syrjanen. Survivin as a marker of cervical intraepithelial neoplasia and high-risk human papillomavirus and a predictor of virus clearance and prognosis in cervical cancer. *Am J Clin Pathol*. 2005. 124:113-21
- 1194 N. Murphy, M. Ring, C. C. Heffron, B. King, A. G. Killalea, C. Hughes, C. M. Martin, E. McGuinness, O. Sheils and J. J. O'Leary. p16INK4A, CDC6, and MCM5: predictive biomarkers in cervical preinvasive neoplasia and cervical cancer. *J Clin Pathol*. 2005. 58:525-34
- 1229 S. Riethdorf, E. F. Neffen, A. Cviko, T. Loning, C. P. Crum and L. Riethdorf. p16INK4A expression as biomarker for HPV 16-related vulvar neoplasias. *Hum Pathol*. 2004. 35:1477-83
- 1247 L. Hu, M. Guo, Z. He, J. Thornton, L. S. McDaniel and M. D. Hughson. Human papillomavirus genotyping and p16INK4a expression in cervical intraepithelial neoplasia of adolescents. *Mod Pathol*. 2005. 18:267-73

- 1250 G. Negri, F. Vittadello, F. Romano, A. Kasal, F. Rivasi, S. Girlando, C. Mian and E. Egarter-Vigl. p16INK4a expression and progression risk of low-grade intraepithelial neoplasia of the cervix uteri. *Virchows Arch.* 2004. 445:616-20
- 1255 M. Branca, M. Ciotti, D. Santini, L. Di Bonito, C. Giorgi, A. Benedetto, P. Paba, C. Favalli, S. Costa, A. Agarossi, M. Alderisio and K. Syrjanen. p16(INK4A) expression is related to grade of cin and high-risk human papillomavirus but does not predict virus clearance after conization or disease outcome. *Int J Gynecol Pathol.* 2004. 23:354-65
- 1270 A. J. Kruse, E. Gudlaugsson, T. Helliesen, E. A. Janssen, B. van Diermen, S. Sandvik and J. P. Baak. Evaluation of prospective, routine application of Ki-67 immunoquantitation in early CIN for assessment of short-term progression risk. *Anal Quant Cytol Histol.* 2004. 26:134-40
- 1276 B. Tringler, C. J. Gup, M. Singh, S. Groshong, A. L. Shroyer, D. E. Heinz and K. R. Shroyer. Evaluation of p16INK4a and pRb expression in cervical squamous and glandular neoplasia. *Hum Pathol.* 2004. 35:689-96
- 1335 G. D. Zielinski, P. J. Snijders, L. Rozendaal, N. F. Daalmeijer, E. K. Risse, F. J. Voorhorst, N. M. Jiwa, H. C. van der Linden, F. A. de Schipper, A. P. Runsink and C. J. Meijer. The presence of high-risk HPV combined with specific p53 and p16INK4a expression patterns points to high-risk HPV as the main causative agent for adenocarcinoma in situ and adenocarcinoma of the cervix. *J Pathol.* 2003. 201:535-43
- 1425 R. Klaes, A. Benner, T. Friedrich, R. Ridder, S. Herrington, D. Jenkins, R. J. Kurman, D. Schmidt, M. Stoler and M. von Knebel Doeberitz. p16INK4a immunohistochemistry improves interobserver agreement in the diagnosis of cervical intraepithelial neoplasia. *Am J Surg Pathol.* 2002. 26:1389-99

- 1480 A. J. Kruse, J. P. Baak, P. C. de Bruin, F. R. van de Goot and N. Kurten. Relationship between the presence of oncogenic HPV DNA assessed by polymerase chain reaction and Ki-67 immunohistochemical features in cervical intraepithelial neoplasia. *J Pathol.* 2001. 195:557-62
- 1507 J. T. Keating, A. Cviko, S. Riethdorf, L. Riethdorf, B. J. Quade, D. Sun, S. Duensing, E. E. Sheets, K. Munger and C. P. Crum. Ki-67, cyclin E, and p16INK4 are complimentary surrogate biomarkers for human papilloma virus-related cervical neoplasia. *Am J Surg Pathol.* 2001. 25:884-91
- 1528 R. Klaes, T. Friedrich, D. Spitkovsky, R. Ridder, W. Rudy, U. Petry, G. Dallenbach-Hellweg, D. Schmidt and M. von Knebel Doeberitz. Overexpression of p16(INK4A) as a specific marker for dysplastic and neoplastic epithelial cells of the cervix uteri. *Int J Cancer.* 2001. 92:276-84
- 1538 A. J. Kruse, J. P. Baak, P. C. de Bruin, M. Jiwa, W. P. Snijders, P. J. Boodt, G. Fons, P. W. Houben and H. S. The. Ki-67 immunohistochemical quantitation in cervical intraepithelial neoplasia (CIN): a sensitive marker for grading. *J Pathol.* 2001. 193:48-54
- 1953 A. Chaux, R. Pfannl, I. M. Rodriguez, J. E. Barreto, E. F. Velazquez, C. Lezcano, A. Piris, G. J. Netto and A. L. Cubilla. Distinctive immunohistochemical profile of penile intraepithelial lesions: a study of 74 cases. *Am J Surg Pathol.* 2011. 35:553-62
- 1955 A. L. Cubilla, B. Lloveras, M. Alejo, O. Clavero, A. Chaux, E. Kasamatsu, N. Monfuleda, S. Tous, L. Alemany, J. Klaustermeier, N. Munoz, W. Quint, S. de Sanjose and F. X. Bosch. Value of p16(INK)(a) in the pathology of invasive penile squamous cell carcinomas: A report of 202 cases. *Am J Surg Pathol.* 2011. 35:253-61

- 1960 A. Chaux, R. Pfannl, B. Lloveras, M. Alejo, O. Clavero, C. Lezcano, N. Munoz, S. de Sanjose, X. Bosch, M. Hernandez-Perez, E. F. Velazquez and A. L. Cubilla. Distinctive association of p16INK4a overexpression with penile intraepithelial neoplasia depicting warty and/or basaloid features: a study of 141 cases evaluating a new nomenclature. *Am J Surg Pathol.* 2010. 34:385-92
- 1962 O. J. Kayes, M. Loddo, N. Patel, P. Patel, S. Minhas, G. Ambler, A. Freeman, A. Wollenschlaeger, D. J. Ralph, K. Stoeber and G. H. Williams. DNA replication licensing factors and aneuploidy are linked to tumor cell cycle state and clinical outcome in penile carcinoma. *Clin Cancer Res.* 2009. 15:7335-44
- 1994 M. Santos, S. Landolfi, A. Olivella, B. Lloveras, J. Klaustermeier, H. Suarez, L. Alos, L. M. Puig-Tintore, E. Campo and J. Ordi. p16 overexpression identifies HPV-positive vulvar squamous cell carcinomas. *Am J Surg Pathol.* 2006. 30:1347-56
- 2111 M. T. Galgano, P. E. Castle, K. A. Atkins, W. K. Brix, S. R. Nassau and M. H. Stoler. Using biomarkers as objective standards in the diagnosis of cervical biopsies. *Am J Surg Pathol.* 2010. 34:1077-87